



**NATIONAL STANDARD OF THE
PEOPLE'S REPUBLIC OF CHINA**

GB 1886.309-2020

National food safety standard - Food additive - Spirulina blue

Issued on: September 11, 2020

Implemented on: March 11, 2021

**Issued by: National Health Commission and State Administration for
Market Regulation**

Table of Contents

1 Scope	3
2 Technical Requirements	3
Appendix A Test Methods	4
Appendix B Ultraviolet-Visible Light Absorption Spectrum of Spirulina Blue	7

1 Scope

This Standard is applicable to the food additive of spirulina blue that uses Spirulina cultivated in fresh water or seawater as raw materials and prepares through water extraction, purification and other processes.

2.1 Sensory requirements

The sensory requirements shall comply with the provisions of Table 1.

Table 1 -- Sensory Requirements

2.2 Physical and chemical indicators

The physical and chemical indicators shall comply with the provisions of Table 2.

Table 2 - Physical and Chemical Indicators

Appendix A Test Methods

A.1 General rules The reagents and water used in this standard refer to analytically-pure reagents and Class-III water specified in GB/T 6682 when no other requirements are indicated. The solution used in the test refers to the aqueous solution when the solvent is not specified.

A.2 Identification test
A.2.1 Solubility It is easily soluble in water; shows blue clear liquid; insoluble in organic solvents such as absolute ethanol, ether, petroleum ether, etc. Take appropriate amounts of spirulina blue specimen (0.05g for powder product; 0.3g for liquid product); dissolve it in 10mL of water; slowly add 4.5g of ammonium sulfate solid powder; mix until it is completely dissolved; and let it stand to form a blue precipitate.

A.2.2 Stability The 0.5% specimen solution prepared by phosphate buffer solution (pH 6.8) shall be blue and accompanied by red fluorescence. After heating at 90°C for 30 min, the fluorescence disappears.

A.2.3 Maximum absorption wavelength
A.2.3.1 Reagents and materials Phosphate buffer:

0.1mol/L, pH 6.8. Prepare as follows:

a) Solution A (0.2 mol/L sodium dihydrogen phosphate aqueous solution): take 31.20g of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$; dissolved and make constant volume by water to 1000mL;

b) Solution B (0.2 mol/L disodium hydrogen phosphate aqueous solution): take 53.61g of $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ (or 71.63g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ or 35.61g of $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$); dissolve and make constant volume by water to 1000mL;

c) Take 51.0mL of solution A and 49.0mL of solution B; and mix evenly and shake well. Add water to dissolve and make the constant volume to 200mL; and obtain 0.1mol/L phosphate buffer solution (pH 6.8).

A.2.3.2 Apparatus