

ICS 65.120
CCS B46



**NATIONAL STANDARD OF THE
PEOPLE'S REPUBLIC OF CHINA**

GB/T 13089-2020

Replacing GB/T 13089-1991

Method for determination of oxazolidinethione in feeds

饲料中噁唑烷硫酮的测定方法

Issued on: November 19, 2020

Implemented on: June 1, 2021

**Issued by: State Administration for Market Regulation;
Standardization Administration of China**

Table of Contents

Foreword...	3
1 Scope...	4
2 Normative References...	4
3 Principle...	4
4 Reagents and Materials...	5
5 Apparatus...	6
6 Samples...	6
7 Test Procedures...	7
8 Test Data Processing...	7
9 Precision...	8

Foreword

This Standard was drafted as per the rules specified in GB/T 1.1-2009. This Standard replaced GB/T 13089-1991 Method for Determination of Oxazolidinethione in Feeds. Compared with GB/T 13089-1991, this Standard has the major differences in technical contents as follows besides the editorial modifications:

--- Modify the scope of application of the method and increase the limit of quantification (see Clause 1 of this Edition; Clause 1 of 1991 Edition);

--- Modify the principle of the method (see Clause 3 of this Edition; Clause 2 of 1991 Edition);

--- Modify the preparation method of white myrosinase (see 4.10 of this Edition;

3.4 of 1991 Edition);

---Modify the dosage and test procedures of white myrosinase (see Clause 7 of this Edition; Clause 6 of 1991 Edition);

--- Modify test data processing (see Clause 8 of this Edition; Clause 7 of 1991 Edition);

--- Modify the precision (see Clause 9 of this Edition; Clause 7 of 1991 Edition). Please note some contents of this document may involve patents. The issuing agency of this document shall not assume the responsibility to identify these patents. This Standard was proposed by and under the jurisdiction of National Technical Committee for Standardization of Feed Industry (SAC/TC 76). Drafting organizations of this Standard: Sichuan Will Test Technology

Co., Ltd.; Institute of Animal Sciences of Chinese Academy of Agricultural Sciences; Public Monitoring Center for Agro-Product of Guangdong Academy of Agricultural Sciences; Tongwei Co., Ltd.; and China Feed Industry Association. Chief drafting staffs of this Standard: Zhang Fengping, Tong Jianming, Wang Weili, Lei Baoliang, Yang Fashu, Wang Liwen, Song Tao, and Lu Jiawen. The historical edition replaced by this Standard is as follows:

--- GB/T 13089-1991. Method for Determination of Oxazolidinethione in Feeds

1 Scope

GB/T 13089-2020 is the Chinese national standard on method for determination of oxazolidinethione in feeds, in the field of agriculture. The /T suffix marks it as a recommended standard: it is not compulsory by itself, but it becomes binding as soon as a contract, a tender or a customer specification calls it up - which in practice is how most foreign buyers meet it. It was issued on 19 November 2020 by the State Administration for Market Regulation; Standardization Administration of China, and has been in force since 1 June 2021. Classification: ICS 65.120, CCS B46. This page is published from the official record of the standard held by the Chinese standards administration: the identification, the dates, the classification and the issuing body are taken from there. The clause text, the tables and the numeric limits are in the document itself, which is delivered complete in English translation.

This Standard specifies the UV spectrophotometric method for the determination of oxazolidinethione in feeds. This Standard is applicable to the determination of oxazolidinethione in rapeseed and its processed products, as well as compound feeds, concentrated feeds and concentrate supplements containing rapeseed and its processed products. The limit of quantification of this Standard method is 150mg/kg.

2 Normative References

The following documents are essential to the application of this document. For the dated documents, only the versions with the dates indicated are applicable to this document; for the undated documents, only the latest version (including all the amendments) is applicable to this document.

GB/T 5520 Inspection of Grain and Oils - Germination Test of Seeds

GB/T 6682 Water for Analytical Laboratory Use - Specification and Test Methods

GB/T 20195 Animal Feeding Stuffs Preparation of Test Samples

3 Principle

The glucosinolate in the specimen is hydrolyzed into isothiocyanate under the action of

white myrosinase in the buffer solution with pH 7.0. After extraction with dichloromethane, the isothiocyanate with hydroxyl reacts with ethanol. The oxazolidinethione is cyclized and formed, which is determined by ultraviolet spectrophotometry.

4 Reagents and Materials

Unless otherwise specified, only use analytically-pure reagents.

4.1 Water: Class-II water specified in GB/T 6682.

4.2 Petroleum ether (boiling range 30°C ~60°C).

4.3 Dichloromethane.

4.4 Absolute ethanol.

4.5 Citric acid solution (0.1mol/L): accurately take 4.20g of citric acid ($C_6H_8O_7 \cdot H_2O$) and dissolve and make constant volume by water to 200mL; and mix evenly. Prepare for current use.

4.6 Disodium hydrogen phosphate (0.2mol/L): accurately take 28.39g of anhydrous disodium hydrogen phosphate (Na_2HPO_4) and dissolve and make constant volume by water to 1L; and mix evenly. Prepare for current use.

4.7 Hydrochloric acid (0.01mol/L): accurately pipette 0.9mL of concentrated hydrochloric acid; dilute and make constant volume by water to 1L; and mix evenly.

4.8 Sodium hydroxide solution (0.01mol/L): accurately take 0.40g of sodium hydroxide and dissolve and make constant volume by water to 1L; and mix evenly.

4.9 Buffer solution with pH 7.0: pipette 176.5 mL of citric acid solution (4.5) in a 1000 mL volumetric flask; dilute and make constant volume by disodium hydrogen phosphate solution (4.6) to 1000 mL; mix evenly. Use hydrochloric acid solution (4.7) or sodium hydroxide solution (4.8) to adjust the pH to 7.0. Prepare for current use.