

ICS 65.100
CCS B 17



**NATIONAL STANDARD OF THE
PEOPLE'S REPUBLIC OF CHINA**

GB/T 17980.29-2026

Replacing GB/T 17980.29-2000

**Pesticides - Guidelines for field efficacy trials - Part 29:
Fungicides against rust of vegetables**

农药 田间药效试验准则 第29部分：杀菌剂防治蔬菜锈病

Issued on: February 27, 2026

Implemented on: September 1, 2026

**Issued by: State Administration for Market Regulation;
Standardization Administration of the PRC**

Table of Contents

- 1 Scope
- 4 Test conditions
- 5 Experimental Design and Arrangement
- 5.2 Reagents
- 5.3 Community Arrangement
- 6 Applying pesticides
- 7 Survey
 - 7.1 Drug Efficacy Survey
 - 7.4 Impacts on other organisms
 - 7.5 Meteorological and Soil Data
- 8 Methods for calculating drug efficacy
 - 8.1 Formula for Calculating Drug Efficacy
- 9 Results and Report Writing

Foreword

This document complies with the provisions of GB/T 1.1-2020 "Standardization Work Guidelines Part

1.Structure and Drafting Rules of Standardization Documents". Drafting. This document is Part 29 of GB/T 17980, "Guidelines for Field Efficacy Testing of Pesticides". GB/T 17980 has already published the following parts.

1.Insecticide control of lepidopteran borers in rice;

2.Insecticide Control of Rice Leaf Roller;

3.Pesticide Control of Rice Leafhoppers; ...

29.Fungicides for the Control of Vegetable Rust; ...

---Part 149: Insecticides for the Control of Red Imported Fire Ants. This document supersedes GB/T 17980.29-2000 "Guidelines for Field Efficacy Tests of Pesticides (I) Control of Vegetable Rust with Fungicides", and is consistent with... Compared with GB/T 17980.29-2000, apart from structural adjustments and editorial changes, the main technical changes are as follows:

a) The scope of application has been changed (see Chapter 1, Chapter 1 of the.2000 edition);

- b) The requirements for test conditions have been changed (see Chapter 4, Chapter 2 of the.2000 edition);
- c) Added requirements for experimental treatments and blank controls (see 5.1);
- d) The requirements for test and control reagents have been changed (see 5.2, 3.1 of the.2000 edition);
- e) Changes were made to cell arrangement, cell size, and duplication requirements (see 5.3, 3.2 of the.2000 edition);

1 Scope

GB/T 17980.29-2026 is the Chinese national standard covering the trial protocol for vegetable rusts - with the pustule counts and severity scales, on diseases that are conspicuous long before they are economically damaging. The GB/T 17980 series is what a pesticide registration in China rests on: it fixes the plot design and replication, the application rate and timing, the assessment of the pest or disease and of the crop, and the analysis of the result. It replaces GB/T 17980.29-2000 and has been in force since 1 September 2026. It was issued on 27 February 2026 and has been in force since 1 September 2026, replacing GB/T 17980.29-2000. The document is under the responsibility of the Ministry of Agriculture and Rural Affairs. This page is published from the official record of the 2026 edition; the clause text of a standard this recent is not yet in circulation, and the figures, limits and tables it contains are those of the document itself, delivered in full with the English translation.

This document describes the method for field plot trials of fungicides for controlling vegetable rust. Rust (*Puccinia allii*), asparagus rust (*Puccinia asparagi*), and broad bean rust (*Uromyces fabae*). [Registered field pesticides for use.] The efficacy of the drug should be evaluated in small-plot trials and other field efficacy trials.

4 Test conditions

4.1 Selection of test subjects, crops and varieties The test subject was rust. The experimental crops were kidney beans, garlic, onions, asparagus, and broad beans, using either susceptible varieties or locally commonly grown varieties. Variety names were recorded.

4.2 Environmental Conditions Field trials should be conducted in disease-prone areas. Cultivation conditions (soil type, fertilization, sowing/planting date, growth stage) for all experimental plots should be considered. The spacing between sections and rows of crops should be uniform and consistent with local scientific agricultural practices. If irrigation is used, record the method, time, and amount of water used. When applying fumigants,

smoke agents, powders (spraying), or fogging agents in greenhouses, each treatment should be performed in a single greenhouse or the greenhouse should be tightly isolated. It is divided into several communities.

5 Experimental Design and Arrangement

5.1 Test Treatment Test reagents, control reagents, and blank controls should be included.

5.2 Reagents

5.2.1 Test reagents The test reagent should be applied at least three times; in special cases, the dosage should be set according to experimental requirements. Specify the Chinese/English common name or code of the reagent. Dosage form, active ingredient content, manufacturer, production date or batch number, expiration date, etc.

5.2.2 Control reagent The control agent should be a registered, locally used product with good efficacy and safety in actual use. The type and function of the control agent should be clearly defined. The method should be the same as or similar to that of the test drug, and the registered dosage should be used. If the test drug is a compound formulation, a single dose of each active ingredient should also be included as a control. Control drug. Special cases may vary depending on the purpose of the experiment. Record the Chinese/English common name, dosage form, active ingredient content, and manufacturer of the control drug. The product name, registration number, production date or batch number, and dosage are required.

5.3 Community Arrangement

5.3.1 Layout of the community The experimental, control, and blank control plots were randomly assigned using a block design, and the plot diagram was recorded. Special circumstances should be noted. illustrate.

5.3.2 Area of the community and duplication Area of the residential area. 15m²~50m² (for sheds, the area should be no less than 8m²). Number of repetitions. at least 4 repetitions.

6 Applying pesticides

6.1 Application Method Follow the test requirements and label instructions. Application methods should be adapted to local scientific agricultural practices. Record the application methods and procedures. process.

6.2 Application equipment Select commonly used equipment in production, or choose equipment according to experimental requirements. Record the type of equipment used and its operating conditions (e.g., working pressure, etc.). All information regarding nozzle type and nozzle diameter.

6.3 Application time and frequency Conduct the experiment according to the test requirements and label instructions. Record the number of applications, the date of each application, and the crop growth stage. The first application should be made at the site of the disease. Before or at the initial stage of an outbreak, the decision to apply pesticide again depends on the development of the disease during crop growth.

6.4 Dosage Administer the medication according to the test requirements and the dosage specified on the label, and record the prepared dosage and the amount of active ingredient. The prepared dosage is usually expressed in mL/667m² (milliliters). The dosage is expressed as per 667 square meters (g/667m²) or g/667m² (grams per 667 square meters), while the effective ingredient dosage is expressed as g/hm² (grams per hectare). Spray application. When applying pesticides, record the amount of pesticide solution used per hectare, expressed in L/hm² (liters per hectare); for fumigants, smoke agents, powders (spraying), and fogging agents. At the same time, record the volume of the greenhouse and the cultivated area, and record the dosage of pesticide per square meter or per cubic meter, in g/m² (grams per square meter) or The dosage is expressed in g/m³ (grams per cubic meter). The pesticide application rate should be evenly distributed, and the deviation should not exceed $\pm 10\%$. If the deviation exceeds $\pm 10\%$, it should be recorded and... Assess its impact.

6.5 Chemical requirements for controlling other diseases, pests, and weeds If other agents are used, those that have no effect on the test agent and the test subjects should be selected, and all plots should be treated uniformly. Furthermore, it should be used separately from the test and control agents to minimize interference from these agents. Accurate records of the application of this type of agent should be kept. Data such as the generic name of the drug, dosage form, content of active ingredient, manufacturer, dosage, application method, application time, and target pests.

7.1 Drug Efficacy Survey

7.1.1 Survey Methodology Five sampling points were taken along the diagonal of each plot. Two plants were surveyed at each point for green beans and broad beans, and ten plants were surveyed for garlic, onions, and asparagus. All leaves of each plant were surveyed. The leaves are graded based on the percentage of the spore mass area on each leaf relative to the total leaf area. Disease investigation and classification are as follows:

0.No disease;

1.The area of the spore mass is less than 5% of the total leaf area;

3.The area of the spore mass accounts for more than or equal to 5% and less than 10% of the total leaf area;

5.The area of the spore mass accounts for more than or equal to 10% and less than 20% of the total leaf area;