

ICS 65.020.40
CCS B 65

LY

**INDUSTRY STANDARD OF THE
PEOPLE'S REPUBLIC OF CHINA**

LY/T 1828-2009

Planting technical criterion of *Pistalia chinensis* Bunge

黄连木栽培技术规程

(English Translation)

Issued on: June 18, 2009

Implemented on: October 1, 2009

Issued by: State Forestry Administration

Table of Contents

Foreword	
1 Scope	
2 Normative references	
3 Seedling selection	
4 Woodland selection	
5 Site preparation	
6 Planting afforestation	
6.1 Planting density	
6.2 Pollination Tree Configuration	
6.3 Planting time	
6.4 Planting methods	
7 Tending management	
7.1 Soil Management	

Foreword

This standard was proposed by the State Forestry Administration.

This standard was drafted. Henan Forestry Technology Extension Station, Linzhou Forestry Technology Extension Station, Anyang Forestry Technology Extension Station, Xinxiang City Forestry Technology Extension Station, Hebi Forestry Technology Extension Station, Huixian Forestry Technology Extension Station.

1 Scope

This standard stipulates the selection of seedlings in the production of *P. chinensis*, the selection of forest land, land preparation, planting and afforestation, tending management, low-yield forest transformation, fruit mining.

Receiving, pest control and other technologies.

This standard applies to the production management of *P. yunnanensis* ecological energy forest planting and afforestation.

2 Normative references

The terms in the following documents become the terms of this standard by reference to this standard. All dated references, followed by all Modifications (not including errata content) or revisions do not apply to this standard, however, parties to agreements based on this standard are encouraged to study it is possible to use the latest version of these files? For undated references, the latest edition applies to this standard.

GB 4285-1989 Standard for safe use of pesticides GB 6000-1999 Quality classification of main afforestation tree seedlings

3 Seedling selection

According to the quality classification of the main afforestation tree species in GB 6000-1999, choose from the seed garden, the mother forest or the fine forest.

High-quality seedlings cultivated on mature mother plants with mature seeds, or improved seedlings cultivated by tissue culture, grafting, cutting and other methods.

4 Woodland selection

According to the local natural conditions, choose a plot with an elevation of 1200m below, a soil thickness of 20cm or more, a leeward sunny, well-drained plot.

Afforestation of berberine.

5 Site preparation

The strip-shaped preparation is 50cm wide and 50cm deep; the hole-shaped preparation is 60cmx60cmx50cm.

6.1 Planting density

The planting density was 825 plants/hm² to 1650 plants/hm².

6.2 Pollination Tree Configuration

The pollination tree is evenly arranged from 1/20 to 1/10.

6.3 Planting time

The bare root seedlings can be planted in the spring before the soil is thawed from March to April, before the seedlings are germinated, and the seedlings are deciduous after autumn and winter. Container seedling it can be planted in the rainy season, spring and autumn.

6.4 Planting methods

The planting depth is preferably buried in the soil to the seedling root line. In arid areas, drought-resistant afforestation techniques such as cutting dry, planting acupoints and covering mulch should be adopted.

7.1.1 Deeply changing soil

In autumn and winter, combined with organic fertilizer, deep turning, expanding holes and changing soil.