Chinastandards.org

SN

INDUSTRY STANDARD OF THE PEOPLE'S REPUBLIC OF CHINA

SN/T 3417-2012

Detection and identification of Carulaspis juniperi(Bouche)

松唐盾蚧检疫鉴定方法

(English Translation)

Issued on: December 12, 2012 **Implemented on: July 1, 2013**

Issued by: General Administration of Customs

Preview: page 1 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3417-2012

Chinastandards.org

Table of Contents

Foreword

1 Scope

2 Terms and definitions

Foreword

This standard was drafted in accordance with GB/T 1.1-2009 given rules.

This standard is proposed and managed by the National Certification and Accreditation Administration Committee.

This standard was drafted. People's Republic of China Shanghai Entry-Exit Inspection and Quarantine Bureau, Inner Mongolia, People's Republic of China Entry-Exit Inspection and Quarantine Bureau, Chinese Academy of Inspection and Quarantine.

1 Scope

This standard defined the plant quarantine Songtang shield Kuwana CarulaspisJuniperi (Bouchè) of quarantine and identification methods.

This standard applies to Songtang shield Kuwana host seedlings, scion, quarantine and identification of fruit plant quarantine Songtang shield Kuwana.

2 Terms and definitions

The following terms and definitions apply to this document.

Tube gland ducts

One kind of glandular structures, have different lengths and diameters of the duct, the duct sides nearly parallel, at the opening of a narrow circle of ossification terrier package Around. According tube gland position can be located in the back of the gland called the penstock pipe gland (dorsalducts), distributed in the body known as the edge of the tube gland edge tube Gland (marginalmacroducts).

Shelly scalecover

Protection structure Diaspididae scale insects, parasites coated on the back of the ventral or dorsal or ventral gland secretions by the parasite hip board, anal gum Excreta and nymphs molt bonded.

Gland plate diskpore Gland plate holes gland, also known as scale insects wax glands

Preview: page 2 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3417-2012

secrete a type of three-hole gland including a variety of shapes, five-hole glands, porous, sieve-like holes, etc.

Glands.

Hip plate pygidium

After the end of the abdominal section number ossification healing from the plate-like body, a shell-forming shield Kuwana related.

Hip leaf pygidiallobes Ossification of the hip edge of the board flat rosette subsidiary projections, in pairs, symmetrically distributed on both sides of the hip edge of the board. Located in the center of the hip plate referred to as A pair hip leaf, also known as the hip leaves; located in a symmetrical on both sides of the second pair of leaf hip hip leaf, later said third and fourth pairs of buttocks leaves.

Hip bar (hip spine) pygidialplates Hip edge of the board is not affiliated ossification flaky projections, a variety of shapes, most of the branches into the comb, sometimes on the buttocks in the previous section, too.

Wai Yin gland perivulvar (disc) pores A disc glands, mostly five grid gland, often 2 to 5 groups located in the vicinity of the vulva Diaspididae female adults.

Hardened rods (thick bars) paraphyses Hip hip board by the edges or sides of the front lobe vertical or inclined elongated body extending hardened structure, sometimes other shapes.

Preview: page 3 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3417-2012