Chinastandards.org

SN

INDUSTRY STANDARD OF THE PEOPLE'S REPUBLIC OF CHINA

SN/T 3574-2013

Detection and identification of Carpomya pardalina(Bigot)

甜瓜实蝇检疫鉴定方法

(English Translation)

Issued on: March 1, 2013 **Implemented on: September 16, 2013**

Issued by: General Administration of Customs

Preview: page 1 of 3 shown, 9 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3574-2013

Chinastandards.org

Table of Contents

Foreword

1 Scope

2 Terms and definitions

Foreword

This standard was drafted in accordance with GB/T 1.1-2009 given rules.

Please note that some of the content of this document may involve patents. Distribution of this document Institutions do not assume the responsibility to identify these patents.

This standard is proposed and managed by the National Certification and Accreditation Administration Committee.

This standard was drafted. People's Republic of China Guangdong Entry-Exit Inspection and Quarantine Zhongshan People's Republic of China Bureau, People's Republic of China Huizhou Entry-Exit Inspection and Quarantine Xishuangbanna People's Republic of China, South China Agricultural the University.

1 Scope

This standard specifies the melon fruit fly [Carpomyapardalina (Bigot)] of quarantine and identification methods.

This standard applies to the quarantine identified melon fruit fly host plants and fruit in the melon fruit fly.

2 Terms and definitions

The following terms and definitions apply to this document (see Appendix A).

Upper forehead bristles frontalbristle Located in the frontal region by the head of the bristles, also known as the amount of bristles.

Amount lower side bristles orbitalbristle Located in the frontal region, and on the number side of the forehead beneath the bristles of bristle, also known as orbital bristles.

Monocular bristles ocelarbristle Located around the bristles monocular.

Cheek bristles genalbristles He was born in cheek bristles.

Inner top bristles innerverticalbristles Monocular near top bristles.

Preview: page 2 of 3 shown, 9 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3574-2013

Outer top bristles outerverticalbristles Near the compound eyes of top bristles.

After eye mane postocularbristles Located behind the trailing edge of the edge of a fly's eye mane, also known as behind columns bristles.

After monocular bristles postocelarbristle Monocular triangle located behind the bristles.

Facial face

In front of the head, between the compound eyes and antennae between area between the mouth of the topsheet.

2.10

Facial spots Located on the face of plaque.

Preview: page 3 of 3 shown, 9 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3574-2013