Chinastandards.org

ICS 07.080
CCS B16

SN

INDUSTRY STANDARD OF THE PEOPLE'S REPUBLIC OF CHINA

SN/T 3765-2013

Detection and identification of Carpomya incompleta(Becker)

欧非枣实蝇检疫鉴定方法

(English Translation)

Issued on: November 6, 2013 **Implemented on: June 1, 2014**

Issued by: General Administration of Customs

Preview: page 1 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3765-2013

Chinastandards.org

Table of Contents

Foreword

1 Scope

2 Terms and definitions

Foreword

This standard was drafted in accordance with GB/T 1.1-2009 given rules.

This standard is proposed and managed by the National Certification and Accreditation Administration Committee.

This standard was drafted. Xinjiang People's Republic of China Entry-Exit Inspection and Quarantine Yili People's Republic of China Bureau of Xinjiang Agricultural University.

1 Scope

This standard specifies the Europe and Africa jujube fruit fly [Carpomyaincompleta (Becker)] of quarantine and identification methods.

This standard applies to the entry and exit of dates and other portable Europe and Africa jujube fruit fly quarantine identification.

2 Terms and definitions

The following terms and definitions apply to this document.

Facial face

In front of the head, between the compound eyes and antennae between area between the mouth of the topsheet.

Facial spots Located on the facial spots. See Appendix A.

Gland spot shiningspots 5 plate one pair of front and slightly sagging gloss region.

Comb hair pectin

1 male adult abdomen after the third quarter gross margin edge of the row.

Side of the caudate lobe surstyli Merging the 9th abdominal segment backplane paired appendages on both sides, from the 10th abdominal segment backplane minutes ago and posterior lobe.

Preview: page 2 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3765-2013

Chinastandards.org

Ovipositor aculeus

Females lay eggs vessel portion for penetrating hole, which often retracted into the base section.

Upper forehead bristles superiorfrontoorbitalsetae Located in the frontal region by the head of the bristles.

Amount lower side bristles inferiorfrontoorbitalsetae it located in the frontal region, and on the number side of the forehead below the mane of bristles.

After monocular mane post-ocelarsetae Monocular triangle located behind the bristles.

2.10

Inner top bristles innerverticalseta Monocular near top bristles.

Preview: page 3 of 3 shown, 11 in the full document. Complete English PDF, instant download:
https://chinastandards.org/standards/sn-t-3765-2013